

Critical Care[™]

DISINFECTANT-FUNGICIDE-VIRUCIDE*



EPA Reg. No. 72977-3-69268

- 30 second⁺ immediate disinfection
- 24-hour[§] residual bacteria kill
- Odorless-no harsh disinfectant smell
- Disinfectant-Fungicide-Virucide
- Reduced toxicity- HMIS Health = 0
- Ready-to-use
- No sticky or dulling residue
- Kills MRSA in 2 minutes

Recommended for use in combination with H₂Orange₂ Concentrate 117[™] for complete, low toxicity cleaning, sanitizing and disinfecting.



30-Second⁺ Kill/ 24-hr.[§] Residual Kill Fast, Easy, Effective

Fastest kill times, 24 hour[§] residual kill, lowest toxicity

Critical Care is formulated to use on critical disease transfer points. Use it in combination with H₂Orange₂ Concentrate 117[™] multi-purpose cleaner-deodorizer-sanitizer, and eliminate elevated toxins from your cleaning and disinfecting program.

⁺ Organisms controlled with 30 second kill time
[§] When used as directed provides protection from *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* up to 24 hours after initial application.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CRITICAL CARE™ is a colorless, odorless broad spectrum antimicrobial disinfectant and deodorizer. Proven to kill bacteria, fungus and viruses*, **CRITICAL CARE** should be used on non-porous environmental hard surfaces in homes, offices, hospitals, nursing homes, medical clinics, dental clinics, pharmacies, laboratories, ambulances and patient transfer vehicles, funeral homes, hotels, restaurants, schools, daycare facilities, veterinary clinics, animal shelters, kennels, cages, stables, catteries, animal transport vehicles, bars, supermarkets, colleges, dorm rooms, churches, basements, garages, workshops, attics, gyms, spas, health clubs, laundromats, school buses, cars, RV's, mobile homes, trucks, trailers, shipping containers, rail cars, cruise ships, exercise facilities, beauty and barber shops, subways, trains, airplanes, ships, buses and other public transportation vehicles, locker rooms, kitchens and restrooms.

CRITICAL CARE™ has been formulated to disinfect hard, non-porous environmental surfaces (painted, glazed tile, plastic, metal, glass, glazed porcelain) and objects including walls, floors, counters, sinks, exterior toilet surfaces, faucet handles, handrails, patio furniture, equipment tables, lab benches, toys, toy boxes, booster chairs, potty seats, laundry hampers, grocery carts, cabinets, tubs, showers, doorknobs, light switch covers, telephones, appliances, stove tops, bed frames, wheelchairs, over-bed tables, examination tables, waste containers, tables, chairs, children's toys, diaper pails, diaper changing tables, bathroom and or kitchen counters, desks, play tables, computer keyboard, jungle gyms, playhouses, child car seat (hard surfaces only), strollers, cribs, playpens, activity centers, tanning beds.

Application Instructions

Pre-clean surfaces prior to using this product.

General Disinfectant: For general disinfection and control of the bacteria *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella enterica*, *Listeria monocytogenes*, Vancomycin resistant *Enterococcus faecium* (VRE), Methicillin resistant *Staphylococcus aureus* (MRSA), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA), Community Associated Methicillin resistant *Staphylococcus aureus* (CA-MRSA-PVL), *Escherichia coli* O157:H7, *Acinetobacter baumannii* and *Campylobacter jejuni* the surface must be completely wet with **CRITICAL CARE™** for 2 minutes. The surface may then be wiped dry with a clean towel. When used as directed, **CRITICAL CARE™** provides residual protection from the *Pseudomonas aeruginosa*, *Staphylococcus aureus*, and *Salmonella enterica* up to 24 hours after initial application. Do not touch treated surface after application if residual protection is to be maintained.

Fungus Control: For effective control of the fungus *Trichophyton mentagrophytes*, the surface must be completely wet with **CRITICAL CARE™** for 10 minutes. The surface may then be wiped dry with a clean towel. Re-apply when cleaning or when new growth appears.

***Viral Control:** To kill Herpes Simplex Type 1 F(1) Strain, Influenza A Virus, Hong Kong strain, Rhinovirus R37 Strain 151-1, Polio Virus Type 2 Lansing Strain the surface must be completely wet with **CRITICAL CARE™** for 10 minutes. The surface may then be wiped with a clean towel.

Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by Influenza A virus. **CRITICAL CARE™** is a broad-spectrum hard surface disinfectant that has been shown to be effective against Influenza A Virus and Avian Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1, formerly called swine flu.

Kills HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings (or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of HIV). Instructions for **Cleaning and Decontamination Against HIV** on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids: **Personal Protection:** When handling items soiled with blood or body fluids, use appropriate barrier protection such as latex gloves, gowns, masks or eye coverings. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this disinfectant. **Contact time:** Apply **CRITICAL CARE™** to area to be treated. The surface must be completely wet with **CRITICAL CARE** for 30 seconds. The surface may then be wiped dry with a clean towel. This contact time will not control all organisms listed on this label. Refer to application instructions for other organisms. **Disposal of Infectious Materials:** Blood and other body fluids should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

Critical Care™ Disinfectant Information

Organism	Kill Time	Organism	Kill Time
* <i>Pseudomonas aeruginosa</i>	30 sec.	<i>Campylobacter jejuni</i>	2 min.
<i>Staphylococcus aureus</i>	2 min.	<i>Trichophyton mentagrophytes</i> (Athlete's Foot Fungus)	10 min.
* <i>Salmonella enterica</i>	30 sec.	**HIV type 1-Strain HTLV IIIB ¹	30 sec.
<i>Listeria monocytogenes</i>	2 min.	*Herpes Simplex Type 1 VR-733 F(1) Strain ²	1 min.
Vancomycin resistant <i>Enterococcus faecium</i> (VRE)	2 min.	*Rotavirus	3 min.
Methicillin resistant <i>Staphylococcus aureus</i> (MRSA)	2 min.	*Human Coronavirus	3 min.
Community Associated Methicillin resistant <i>Staphylococcus aureus</i> (CA-MRSA)	2 min.	*Norovirus	10 min.
Community Associated Methicillin resistant <i>Staphylococcus aureus</i> (CA-MRSA-PVL)	2 min.	*Influenza A VR-544, Hong Kong Strain	10 min.
<i>Escherichia coli</i> O157:H7	2 min.	*Rhinovirus R37 VR-1147, Strain 151-1	10 min.
<i>Acinetobacter baumannii</i>	2 min.	*Polio Type 2, VR-1002 Lansing Strain	10 min.

Toxicology & Safety Information

Skin irritation = non irritant	Flammability = non combustible
Eye irritation = non irritant	Dot Classification = non regulated
Acute Oral Toxicity = >5000 mg/kg	Hazardous waste = no
Dermal Sensitization = not a sensitizer	

⁺ Organisms controlled with 30 second kill time.

^{*} When used as directed Critical Care kills HIV type 1-Strain HTLV IIIB, Herpes Simplex Type 1 VR-733 F(1) Strain, Influenza A VR-544, Hong Kong Strain, Rhinovirus R37 VR-1147, Strain 151-1, Polio Type 2, VR-1002 Lansing Strain, Rotavirus, Norovirus, Human Coronavirus and Avian Influenza A.

[§] When used as directed provides protection from *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* up to 24 hrs after initial application.